

PRESENT STATUS OF ELECTROMAGNETIC REACTIONS ON THE DEUTERON ABOVE PION THRESHOLD*

H. ARENHÖVEL, E.M. DARWISH[†], A. FIX, M. SCHWAMB
*Institut für Kernphysik, Johannes Gutenberg-Universität
 55099 Mainz, Germany*

Received (received date)

Revised (revised date)

The present status of the theoretical description of electromagnetic reactions on the deuteron above pion threshold is reviewed. Three major topics are considered: (i) retardation effects in π -meson exchange contributions to NN -interaction and meson exchange currents in deuteron photodisintegration above π -threshold in the Δ -resonance region, (ii) off-shell effects in the one-body current treated in a simple pion cloud model in deuteron photodisintegration in and above the Δ -resonance region, and (iii) final state interaction effects in photoproduction of π and η mesons on the deuteron.

1. Introduction

In view of the fact that at present QCD in the non-perturbative regime can at best be described by effective degrees of freedom only, one uses a framework with meson, nucleon and isobar degrees of freedom. Hadron properties are either described phenomenologically or by effective quark models. The central question then is: How accurate is this effective description, and where is the borderline beyond which explicit quark-gluon degrees of freedom have to be considered? It is very likely that no clear cut answer exists.

However, the study of electromagnetic reactions on few-nucleon systems may give at least a partial answer because lightest nuclei (deuteron, helium-3) allow reliable theoretical descriptions, and approximations, which are unavoidable in more complex many-body systems, are not necessary. Therefore, such systems constitute reliable test laboratories for the investigation of effective degrees of freedom. Furthermore, the electromagnetic interaction is well known and sufficiently weak, in order to allow conclusive interpretations in terms of charge and current matrix elements. Finally, reactions above pion threshold are of particular interest with respect to explicit meson degrees of freedom and internal baryon structure.

The main points of interest are (i) the role of meson and isobar degrees of freedom in medium energy reactions, (ii) many-body phenomena, induced by the effective description in terms of meson, nucleon and isobar d.o.f., e.g., the role of pion retardation in NN interaction and two-body meson exchange operators, and (iii) properties of the neutron like, e.g., the elementary π and η photoproduction on the neutron, in other words, the use of light nuclei as effective neutron targets. Particularly suited are quasifree reactions on the deuteron in order to minimize final

*Supported by Deutsche Forschungsgemeinschaft (SFB 443).

[†]Supported by Deutscher Akademischer Austauschdienst. Present address: Physics Department, Faculty of Science, South Valley University, Sohag, Egypt.

state interaction effects. However, for a reliable interpretation it is mandatory to correct for medium influences as, e.g., described by two-body effects.

2. Pion Retardation in $d(\gamma, p)n$ above Pion Threshold

Most sophisticated theories of photodisintegration in the Δ region, which are based on a coupled channel approach and use a Δ excitation operator from a fit of π photoproduction on the nucleon, encounter various problems¹ (see Fig. 1): (i) Under-




Fig. 1. Total and differential cross sections, and photon asymmetry of deuteron photodisintegration. Dashed: static calculation of Wilhelm et al.¹. Exp.: R. Crawford *et al.*, *Nucl. Phys. A* **603**, 303 (1996), LEGS 95: G. Blanpied *et al.*, *Phys. Rev. C* **52**, R455 (1995), F. Adamian *et al.*, *J. Phys. G* **17**, 1189 (1991), DAPHNE 97: S. Wartenberg *et al.*, *Few-Body Syst.* **26**, 213 (1999).




Fig. 2. Total cross section for $d(\gamma, p)n$. Dashed: static calculation of Wilhelm et al.¹; all other curves from²: dotted: improved static calculation using Bonn OBEPR potential; dash-dot: retardation in hadronic part but static MEC; solid: complete calculation including πd -channel and $\rho\pi\gamma/\omega\pi\gamma$ -MECs. Exp. as in Fig. 1.

estimation of the total cross section by 20-30%. (ii) Angular distributions develop a dip around 90 degree, especially above 300 MeV. (iii) The shape of the photon asymmetry is at variance with experimental data. A detailed analysis of the role of the Born terms and their correspondence to MEC has lead to the conjecture, that π retardation in potential and exchange current might be important. Indeed, in a recent calculation^{2,3} the importance of retardation in both, the NN -interaction as well as MEC has been shown. For the total cross section this is demonstrated in Fig. 2 where a quantitative agreement with experimental data is achieved. Also for the differential cross section and the photon asymmetry shown in Fig. 3 one notes a much improved description. In particular, the dip in the angular distribution has




Fig. 3. Differential cross sections (upper panels) and photon asymmetries (lower panels) of deuteron photodisintegration from². Notation as in Fig. 2. Exp. as in Fig. 1.

disappeared completely. Thus any realistic description of e.m. reactions on light nuclei in this energy regime has to use retarded interactions and MEC.

3. Off-shell Effects in One-Body Current

It is a well-known fact that the e.m. current of a particle with internal structure becomes in general more complicated in an off-shell situation. Consider, for example the e.m. current of a proton. Its on-shell Dirac current is determined by two structure functions, the Dirac and Pauli form factors $F_1(q^2)$ and $F_2(q^2)$, respectively, which depend on q^2 , the four momentum transfer squared, only. However, for e.m. reactions on nuclei the interacting nucleons are off-shell, i.e. $p'^2 = W'^2 \neq M^2 \neq W^2 = p^2$. In this case, one has ten additional structure




Fig. 4. Electromagnetic current of a nucleon dressed with a meson cloud.

functions depending not only on q^2 alone but also on the off-shell masses W' and W . The problem is that for the off-shell structure functions one needs a dynamical model for the internal structure, because these form factors are intimately connected to the underlying hadronic interaction. Thus, such a model has to be consistent with the NN -interaction. Moreover, off-shell effects as such are not independently observable⁴. Recently, we have completed a study of off-shell effects in deuteron photodisintegration⁵ using a consistent dynamical approach in which the nucleon is dressed by pion loops, i.e. the internal nucleon structure is described by a pion




Fig. 5. Differential cross sections. Dashed curves without and solid with off-shell effects from⁵. Exp.: black circles: Crawford *et al.*, NPA **603**, 303 (1996), open squares: Arends *et al.*, NPA **412**, 509 (1984), open circles: G. Blanpied *et al.*, *Phys. Rev. C* **52**, R455 (1995).

cloud. Correspondingly, one finds for the e.m. one-nucleon current the representation depicted in Fig. 4. For the resulting off-shell one-nucleon current one finds in the nonrelativistic limit

$$\langle \vec{p}', e' | \vec{j}(0) | \vec{p}, e \rangle = 2\beta(e', e)\vec{p}' + \gamma(e', e)\vec{k} + i\vec{\sigma} \times (\delta(e', e)\vec{p}' + \epsilon(e', e)\vec{k}),$$

with $\vec{k} = \vec{p}' - \vec{p}$, and e' and e denote final and initial nucleon energies, respectively. The structure functions fulfil the on-shell condition, i.e. for $e' = p'^2/2M$ and $e = p^2/2M$

$$\beta(e', e) = -\gamma(e', e) = \frac{1}{2M}, \quad \delta(e', e) = 0, \quad \epsilon(e', e) = \frac{\mu}{2M}.$$

The correct on-shell current is ensured by an appropriate counter term. Within this meson-nucleon model, which is also used for the NN -interaction including retardation effects, one finds a sizeable influence from off-shell effects on the differential cross section of deuteron photodisintegration in and above the region of the Δ -resonance as is seen in Fig. 5. They show up predominantly at forward and backward angles, leading to a decrease of the cross section. In the future, more realistic nucleon models should be studied with respect to such off-shell effects.

4. Final State Interaction in Incoherent Pion Photoproduction

Meson photoproduction on the nucleon provides valuable information on its internal structure and serves as a test of hadron models. The production on light nuclei is of particular interest because it allows one to study the elementary neutron amplitude, medium effects and nuclear structure. Recently, we have completed a calculation of pion photoproduction on the deuteron⁶ including besides the impulse approximation complete rescattering in all two-body subsystems as depicted in Fig. 6. Deuteron




Fig. 6. Diagrams for incoherent pion photoproduction on the deuteron: (a) impulse approximation (IA), (b) NN rescattering, (c) πN rescattering.




Fig. 7. Total cross section for pion photoproduction on the deuteron from⁶. Dashed: IA; solid: IA+complete rescattering; data: Asai *et al.*, *Phys. Rev. C* **42**, 837 (1990), Benz *et al.*, *Nucl. Phys. A* **65**, 158 (1973) (ABHHM), Chiefari *et al.*, *Lett. Nuovo Cim.* **13**, 129 (1975) (Frascati).




Fig. 8. Differential cross sections for $d(\gamma, \pi^0)pn$ from⁶. Dashed: IA; solid: IA+complete rescattering; Exp.: Krusche *et al.*, *Eur. Phys. J. A* **6**, 309 (1999).

wave function and NN interaction are taken from the Paris potential, and the NN and πN interactions in separable form. The complete T -matrices are obtained from solutions of the corresponding LS-equations. For NN rescattering all partial waves with $J \leq 3$, and for πN rescattering S through D waves are included. Total cross sections are shown in Fig. 7. For charged pion production the rescattering effect is small, but for neutral pion production quite sizeable, which mainly stems from the fact that in IA quite a fraction of the coherent production is included due to the non-orthogonality of the final plane wave to the deuteron bound state. In all cases πN -rescattering is very small compared to NN -rescattering due to the much weaker πN -interaction. The inclusion of such rescattering contributions leads to a satisfactory description of the experimental total cross sections for π^- as well as for π^0 production as shown in Fig. 7. A corresponding good agreement is achieved for the differential cross sections of π^0 production depicted in Fig. 8. One readily notices that the major rescattering effects appear at forward meson angles. This is also true for charged pion production⁶.

5. Two-Body Effects in Coherent Eta Photoproduction

Photoproduction of η mesons is an interesting tool for studying the ($s = 1/2$)-




Fig. 9. Left panel: Various mechanisms for η production on the deuteron. Right panel: Treatment of different time orderings of hadronic transition potentials. (a) static approximation for $NN \leftrightarrow NR$ potential, (b) upper part: retarded NR exchange potential, lower part: static approximation of the meson- RR propagator.




Fig. 10. Differential and total cross section for coherent η -production on deuteron. Left panel: short-dashed: IA, long-dashed: IA + NN resc. + static NS_{11} -resc.; dash-dotted: IA + NN resc. + retarded NS_{11} -resc.; solid: IA + all retarded rescatterings but no MEC. Right panel: dotted: pure resonance contribution; long-dashed: IA, short-dashed: IA + retarded first order rescattering, solid: complete calculation.

nucleon resonances. The $S_{11}(1535)$ plays a special role because of its strong coupling to the η channel. While the incoherent reactions yields estimates of the modulus of the neutron amplitude, the coherent process provides the phase information. Furthermore, the deuteron with $T = 0$ serves as isospin filter and thus yields the ratio of isoscalar to proton amplitude A_s/A_p . Until recently, a seeming discrepancy was noted between extraction from the coherent reaction, $|A_s/A_p|_{coh} \approx 0.2$, and from the incoherent one, $|A_s/A_p|_{incoh} = 0.09$. The latter value was extracted from the incoherent production on the deuteron, yielding at resonance $(\sigma_n/\sigma_p)_{res} = 0.66$.

In recent work on coherent photoproduction on the deuteron⁷ we have taken for the elementary production amplitude a coupled channel model of⁸, which considers the channels $\pi N \rightarrow \pi N$, $\pi N \rightarrow \eta N$, $\gamma N \rightarrow \pi N$, and $\gamma N \rightarrow \eta N$. Specific features

of the model are the inclusion of self energy contributions from π and η loops and the dressing of the e.m. vertex by hadronic rescattering. A good description of




Fig. 11. Left panel: Differential cross section: dashed: IA, solid: complete calculation; data: Hoffmann-Rothe *et al.*, *Phys. Rev. Lett.* **78**, 4697 (1997). Right panel: Linear photon asymmetry: short-dashed: pure resonant contribution, long-dashed: IA, dotted: IA + retarded rescattering, solid: complete calculation.

η photoproduction on the proton is achieved with the important result that the dressing leads to complex values for proton and neutron amplitudes, i.e., $A_n = (-114 - i1.7) \times 10^{-3} \text{GeV}^{-1/2}$, $A_p = (120.9 - i66.1) \times 10^{-3} \text{GeV}^{-1/2}$, yielding the ratios $(\sigma_n/\sigma_p)_{res} = |A_n/A_p|^2 = 0.68 \approx 2/3$, and $A_s/A_p = 0.25 e^{-i0.969}$. Thus there is no contradiction for this ratio anymore between the extraction of this ratio from the coherent and the incoherent reaction.

In the left panel of Fig. 9 the various mechanisms, taken into account in⁷, are displayed. The box labeled Born contains disconnected diagrams where the photon is absorbed by one nucleon and the η is emitted by the other. Hadronic rescattering is indicated by boxes T_{NN} , T_{NR} , T_{RN} , and T_{RR} and meson exchange current contributions by boxes $N[2]$. As resonances “R” we have included $P_{11}(1440)$, $S_{11}(1535)$ and $D_{13}(1520)$. The hadronic interaction is considered in static as well as retarded form as displayed in the right panel of Fig. 9. For coherent η photoproduction on the deuteron, the effect of various mechanisms on the differential cross section are displayed in the left panel of Fig. 10. One notes an opposite effect between static and retarded rescattering. The total cross section is shown in the right panel of Fig. 10, where a sizeable influence from all two-body effects is seen. Furthermore, the first order rescattering approximation overestimates considerably the total cross section. For the differential cross section a reasonable though not perfect agreement with experiment is achieved as is shown in the left panel of Fig. 11. In the right panel of this figure the photon asymmetry is displayed. One notes a sizeable influence from hadronic rescattering but little effect from MEC.

6. Final State Interaction in Incoherent Eta Photoproduction

Near threshold the impulse approximation yields a very small cross section for $d(\gamma, \eta)np$ due to the large momentum mismatch and indeed fails drastically compared to experiment yielding a cross section much too low.




Fig. 12. Left panel: Total cross section for $d(\gamma, \eta)np$ from⁹; dashed: IA, solid: IA + rescattering, dash-dotted: IA + NN rescattering, data: inclusive $\gamma d \rightarrow \eta X$ from¹¹. Middle and right panels: η -meson spectra at forward emission angles for two different photon energies and angles: dotted: IA, solid: IA + NN rescattering, dashed: without the deuteron D -wave in the NN -rescattering contribution. The excitation energy E_{np} in the final NN -system is indicated at the top abscissa.




Fig. 13. Differential cross sections near threshold from⁹. Dotted: IA, dashed: IA + NN rescattering; solid: IA + complete rescattering. Exp. from Krusche *et al.*, *Phys. Lett. B* **358**, 40 (1995).




Fig. 14. η meson spectrum for $d(\gamma, \eta)np$. Dotted: IA, dashed: first order rescattering, solid: complete three-body calculation.

Therefore, we first have performed an approximate treatment of FSI⁹ in complete analogy to pion photoproduction on the deuteron, i.e. taking into account only complete rescattering in the two-body ηN and NN subsystems in the final state. In the following, this is called first order rescattering. In this case the NN t -matrix is determined from the Bonn OBEPQ potential and for the ηN t -matrix an isobar model is taken describing the intermediate $S_{11}(1535)$ excitation. The first order rescattering, restricted to s -waves of NN and ηN subsystems in view of the near-

threshold region, leads to a considerable improvement (see left panel of Fig. 12). The spectrum of the outgoing η meson (middle and right panels of Fig. 12) shows the distinct signature of the final state NN rescattering exhibiting the prominent 1S_0 peak near the NN scattering threshold. The differential cross sections near




Fig. 15. Results for $d(\gamma, \eta)np$ from¹⁰. Left panel: Total cross section: (a) incoherent: dotted: IA; dashed: first order rescattering; dash-dot: complete 3-body model; (b) inclusive: solid: sum of coherent and incoherent channels. Inclusive data (Mainz) from Hejny *et al.*¹². Middle and right panels: Differential cross sections: dotted: IA; dashed: first order rescattering; solid: complete 3-body model. Inclusive data from Hejny *et al.*¹².

threshold are shown in Fig. 13. The left panel of Fig. 13 indicates that first order rescattering still fails to explain quantitatively the enhancement of the data right above threshold. This is corroborated by very recent more precise near-threshold data of Hejny *et al.*¹².

For this reason, we then have performed a three-body treatment of the final state interaction¹⁰, because the very strong effect in first order rescattering suggests that a genuine three-body treatment is required. A considerable simplification is achieved by restriction to only s -waves which is justified because of threshold region. For the NN interaction a simple Yamaguchi form is used. The resulting η spectrum is displayed in Fig. 14 where again one notes clearly the 1S_0 peak as in Fig. 12. However, for the lower photon energy one readily sees a substantial underestimation of the first order rescattering compared to the three-body calculation, although the forms are similar. Total and differential cross sections are shown in Fig. 15. The inclusive total cross section data exhibit a distinct enhancement near threshold which is reproduced by the 3-body approach (left panel of Fig. 15) but not in first order, the latter being considerably lower right at threshold. This is also the case for the differential cross sections (middle and right panels of Fig. 15). It remains to be seen, whether a more realistic treatment of the NN -interaction is also able to describe the data.

7. Conclusions and outlook

In summary, we may conclude that the electromagnetic probe is a very important tool in order to reveal the internal structure of hadrons. Only the new generation of high duty cycle machines allows one to exploit its full power and the thrust of future experimental research lies on the study of exclusive reactions. Polarization observables will give us much more detailed information and thus will provide much more stringent tests for theoretical models.

Reactions on the deuteron are of particular importance for testing present theoretical frameworks for describing strong interaction physics in terms of effective degrees of freedom, thus serving as a test laboratory. Of special interest are e.m. reactions above the pion threshold. An important example is photodisintegration with respect to the study of retardation and off-shell effects.

Furthermore, meson production on the deuteron offers the possibility to study the elementary production amplitude on the neutron provided one has the two-body effects from final state interaction and e.m. current under control. In particular, the ηN -interaction can be studied in incoherent eta production near threshold. However, a first order rescattering calculation as used, e.g. in¹³, is not reliable for that purpose, because right above threshold a three-body approach is mandatory.

Finally, for increased energy and momentum transfers, the effects which arise from relativity should be carefully considered.

1. P. Wilhelm and H. Arenhövel, *Phys. Lett. B* **318**, 410 (1993).
2. M. Schwamb, H. Arenhövel, and P. Wilhelm, *Phys. Lett. B* **420**, 255 (1998).
3. M. Schwamb and H. Arenhövel, *Nucl. Phys. A* **690**, 682 (2001).
4. S. Scherer and H. Fearing, *Nucl. Phys. A* **684**, 499 (2001).
5. M. Schwamb and H. Arenhövel, *Nucl. Phys. A* **696**, 556 (2001).
6. E.M. Darwish, H. Arenhövel, and M. Schwamb, nucl-th/0208030; *Eur. Phys. J. A* (in print).
7. F. Ritz and H. Arenhövel, *Phys. Rev. C* **64**, 034005 (2001).
8. C. Bennhold and H. Tanabe, *Nucl. Phys. A* **530**, 625 (1991).
9. A. Fix and H. Arenhövel, *Z. Phys. A* **359**, 427 (1997).
10. A. Fix and H. Arenhövel, *Nucl. Phys. A* **697**, 277 (2002).
11. B. Krusche *et al.*, *Phys. Lett. B* **358**, 40 (1995).
12. V. Hejny *et al.*, *Eur. Phys. J. A* **13**, 493 (2002).
13. A. Sibirtsev *et al.*, *Phys. Rev. C* **65**, 044007 (2002).